


Weight Loss Surgery

Losing 41% to 66%¹⁻³ of excess weight may be just the beginning...



- Migraines
46% improved⁴
- Depression
47% reduced⁵
- Pseudotumor cerebri
96% resolution of headaches⁶
95% resolution of pulsatile tinnitus⁶
- Obstructive sleep apnea
45% to 76% resolved^{7,8}
- High cholesterol
71% to 94% improved^{9*}
- Asthma
39% resolved¹⁰
- High blood pressure
42% to 66% resolved^{7,8,11}
- Nonalcoholic fatty liver disease
37% resolution of steatosis¹²
- Metabolic syndrome
80% resolved¹²
- GERD
72% to 95% resolved^{5,13}
- Type 2 diabetes
45% to 68% resolved^{7,14}
- Polycystic ovarian syndrome
52% resolution of hirsutism¹⁵
100% resolution of menstrual dysfunction¹⁵
- Urinary stress incontinence
50% resolved¹⁶
- Osteoarthritis/degenerative joint disease
41% resolved⁵
- Venous stasis disease
95% resolution of venous stasis ulcers¹⁷

There are different types of bariatric surgery procedures, and your outcomes may vary depending on the type of procedure (as indicated in the ranges provided above). Please talk with your doctor about potential surgical outcomes related to your procedure.

Ask your doctor about weight loss surgery as a treatment option

or go to www.REALIZE.com

IMPORTANT SAFETY INFORMATION: Bariatric surgery is used in morbidly obese adult patients for significant long-term weight loss. Bariatric surgery may not be right for individuals with certain digestive tract conditions. All surgery presents risks. Weight, age and medical history determine your specific risk. Ask your doctor if bariatric surgery is right for you.

Resolution observed in the context of studies. EES has no independent data to suggest permanent resolution.
*Figure is for hyperlipidemia. Hyperlipidemia is a general term for high fats in blood, which may include cholesterol and / or triglycerides.

1. Phillips E, Ponce J, Cunneen SA, et al. Safety and effectiveness of REALIZE® adjustable gastric band: 3-year prospective study in the United States. *Surg Obes Rel Dis.* 2009; 5:588-597. 2. Fischer L, Hildebrandt C, Bruckner T, Kenngott H, Linke GR, Gehrig T, Büchler MW, Müller-Stich BP. Excessive weight loss after sleeve gastrectomy: a systematic review. *Obes Surg.* 2012 May;22(5):721-31. 3. O'Brien PE, McPhail T, Chaston TB, et al. Systematic review of medium-term weight loss after bariatric operations. *Obes Surg.* 2006; 16(8):1032-1040. 4. Bond DS, Vithianathan S, Nash JM, et al. Improvement of migraine headaches in severely obese patients after bariatric surgery. *Neurology.* 2011 Mar 29;76(13):1135-8. 5. Schauer PR, Ikramuddin S, Gourash W, et al. Outcomes after laparoscopic Roux-en-Y gastric bypass for morbid obesity. *Ann Surg.* 2000 Oct;232(4):515-29. 6. Sugerman HJ, Felton WL 3rd, Sismanis A, et al. Gastric surgery for pseudotumor cerebri associated with severe obesity. *Ann Surg.* 1999 May;229(5):634-40; discussion 640-2. 7. Tice JA, Karliner L, Walsh J, et al. Gastric banding or bypass? A systematic review comparing the two most popular bariatric procedures. *Am J Med.* 2008 Oct;121(10):885-93. 8. EES weighted analysis of data summarized in table 4 of Brethauer SA, Hammel JP, Schauer PR. Systematic review of sleeve gastrectomy as staging and primary bariatric procedure. *Surg Obes Rel Dis.* 2009; 5:469-475. 9. Buchwald H, Avidor Y, Braunwald E, et al. Bariatric surgery: a systematic review and meta-analysis. *JAMA.* 2004 Oct 13;292(14):1724-37. 10. Reddy RC, Baptist AP, Fan Z, et al. The effects of bariatric surgery on asthma severity. *Obes Surg.* 2011 Feb;21(2):200-6. 11. EES analysis of data from US Clinical Trial PMA 070009. 12. Mattar SG, Velcu LM, Rabinovitz M, et al. Surgically-induced weight loss significantly improves nonalcoholic fatty liver disease and the metabolic syndrome. *Ann Surg.* 2005 Oct;242(4):610-7. 13. DeMaria EJ, Sugerman HJ, Kellum JM, et al. Results of 281 consecutive total laparoscopic Roux-en-Y gastric bypasses to treat morbid obesity. *Ann Surg.* 2002 May;235(5):640-5. 14. Based on STAMPEDE population which consisted of predominately female, moderately obese patients with long standing and uncontrolled T2DM. 45% to 68% achieved partial or complete remission of diabetes (diabetes resolution based on HbA1c<7.0). Schauer PR, Kashyap SR, Wolski K, et al. Bariatric surgery versus intensive medical therapy in obese patients with diabetes. *N Engl J Med.* 2012 Apr 26;366(17):1567-76. 15. Eid GM, Cottam DR, Velcu LM, et al. Effective treatment of polycystic ovarian syndrome with Roux-en-Y gastric bypass. *Surg Obes Relat Dis.* 2005 Mar-Apr;1(2):77-80. 16. Kuruba R, Almahmeed T, Martinez F, et al. Bariatric surgery improves urinary incontinence in morbidly obese individuals. *Surg Obes Relat Dis.* 2007 Nov-Dec;3(6):586-90. 17. Sugerman HJ, Sugerman EL, Wolfe L, et al. Risks and benefits of gastric bypass in morbidly obese patients with severe venous stasis disease. *Ann Surg.* 2001 Jul;234(1):41-6.